

Acute Pyelonephritis

Acute pyelonephritis (APN) is a bacterial infection of the kidneys. Imaging tests are usually not recommended for individuals suspected to have APN and without histories of pyelonephritis, diabetes, immune compromise, stones, kidney obstruction, prior kidney surgery, vesicoureteral reflux, lack of response to prior treatment, pregnancy, or advanced age.

For individuals with prior histories listed above, CT abdomen and pelvis (<https://www.radiologyinfo.org/en/info/abdominect>) with contrast is usually appropriate. Ultrasound (US) abdomen (<https://www.radiologyinfo.org/en/info/abdominus>) ; US color Doppler of the kidneys (<https://www.radiologyinfo.org/en/info/ultrasound-renal>) , bladder, and retroperitoneum; MRI abdomen/pelvis (<https://www.radiologyinfo.org/en/info/mri-abdomen-pelvis>) without contrast; MRI abdomen without and with contrast; or CT abdomen/pelvis without contrast may be appropriate.

For individuals with history of kidney stones or kidney obstruction, CT abdomen/pelvis with contrast or CT abdomen/pelvis without and with contrast is usually appropriate. US abdomen; US color Doppler kidneys, bladder, and retroperitoneum; MRI abdomen and pelvis without contrast; MRI abdomen without and with contrast; CT abdomen/pelvis without contrast; or CT abdomen/pelvis without and with contrast may be appropriate.

For pregnant women without complications, US abdomen; US color Doppler of kidneys, bladder, and retroperitoneum; MRI abdomen/pelvis without contrast; MRI abdomen without contrast; or MRU (<https://www.radiologyinfo.org/en/info/urography>) without contrast may be appropriate.

For individuals with a kidney transplant and without removal of native kidneys and without other complications, US duplex Doppler of the kidney transplant or CT abdomen/pelvis with contrast is usually appropriate. MRI abdomen/pelvis without and with contrast, MRI abdomen/pelvis without and with contrast, MRI abdomen/pelvis without contrast, CT abdomen/pelvis without contrast, or CT abdomen/pelvis without and with contrast may be appropriate.

— By Elaine Liang and Aya Kamaya, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple

distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2025 Radiological Society of North America, Inc.