

Supplemental Breast Cancer Screening Based on Breast Density

Breast cancer (<https://www.radiologyinfo.org/en/info/breast-cancer>) is the most common cancer among US women except for nonmelanoma skin cancer.

Breast tissue density can range from nondense breasts that are mostly fatty to dense breasts (<https://www.radiologyinfo.org/en/info/dense-breasts>) that are fibrous and have more glandular tissue. Mammography (<https://www.radiologyinfo.org/en/info/mammo>) is best used to find cancer in nondense breasts. Patients with dense breast tissue may need additional tests.

For women with an average risk for breast cancer (less than 15% lifetime risk) with nondense breasts, digital breast tomosynthesis (DBT) (<https://www.radiologyinfo.org/en/info/tomosynthesis>) screening, also called 3-D mammography, is usually appropriate.

For women with an intermediate risk for breast cancer (15%-20% lifetime risk) with nondense breasts, DBT is usually appropriate, and MRI (<https://www.radiologyinfo.org/en/info/breastmr>) with and without intravenous (IV) contrast as well as abbreviated MRI with and without contrast (shorter MRI) may also be appropriate.

For women with a high risk for breast cancer (greater than 20% lifetime risk) with nondense breasts, DBT and MRI with and without IV contrast are usually appropriate. Mammography with contrast, whole-breast ultrasound (<https://www.radiologyinfo.org/en/info/breastus>), and abbreviated MRI with and without contrast may also be appropriate.

For women at average risk with dense breasts, DBT is usually appropriate. Mammography with contrast and MRI with and without contrast may be appropriate.

For women at intermediate risk with dense breasts, DBT is usually appropriate. Mammography with contrast, breast ultrasound, and MRI with and without contrast may be appropriate.

For women at high risk with dense breasts, DBT, breast ultrasound, and MRI with and without contrast (including abbreviated) are usually appropriate. Mammography with contrast may be appropriate.

For more information, please see the Breast Cancer Screening (<https://www.radiologyinfo.org/en/info/screening-breast>) and Dense Breasts (<https://www.radiologyinfo.org/en/info/dense-breasts>) pages.

— By Leslie Feldman and Kemi Babagbemi, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

Copyright © 2025 Radiological Society of North America, Inc.