

## Breast Implant Evaluation

Breast implants are made of saline, silicone, or both. Some complications that may occur with implants include the implant bursting (rupture), unexplained swollen lymph nodes in the armpits, or a rare blood cancer called anaplastic large-cell lymphoma (<https://www.radiologyinfo.org/en/info/lymphoma>) that may occur a year after surgery. Patients with implants and without symptoms should continue routine breast screening (<https://www.radiologyinfo.org/en/info/screening-breast>).

A saline-implant rupture is often diagnosed with a physical examination. If imaging is needed, ultrasound (<https://www.radiologyinfo.org/en/info/breastus>) (US) should be used for people under 30. For people 30 to 39 years old, a mammogram (<https://www.radiologyinfo.org/en/info/mammo>), digital breast tomosynthesis (<https://www.radiologyinfo.org/en/info/tomosynthesis>) (DBT), or US may be used. A mammogram or DBT is the best test for people 40 and older.

MRI (<https://www.radiologyinfo.org/en/info/breastmr>) without contrast is the best test for diagnosing a silicone-implant rupture, especially when the silicone is still contained by the outer shell (majority of ruptures). Mammography and DBT are not recommended for contained ruptures.

For ruptures outside the shell, in addition to MRI without contrast, US may be used in people under the age of 30. Mammography, DBT, and US may be used for people over 30. For people with prior silicone implants, results may show leftover silicone, not a rupture of new implants, making it important to compare with prior tests.

For unexplained swollen lymph nodes in the armpits, US should be used for patients under the age of 30. For people over 30, mammography, DBT, or US should be used. If the person is suspected to have anaplastic large-cell lymphoma, US is recommended.

— By Celena Romero, RD, MBA, CPHQ, and Ryan K. Lee, MD, MBA, MRMD. This information originally appeared in the *Journal of the American College of Radiology*.

### Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

**Note:** Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

### Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

