

Renal Failure

Renal (kidney) failure (<https://www.radiologyinfo.org/en/info/kidneyfailure>) occurs when one or both kidneys do not function well. Acute kidney failure happens suddenly and can be temporary or progress to chronic failure. Chronic kidney failure is persistent and can slowly worsen over time. In general, contrast should be avoided in kidney failure, unless medically necessary.

For acute kidney injury, kidneys retroperitoneal ultrasound (<https://www.radiologyinfo.org/en/info/ultrasound-renal>) is usually appropriate (ultrasound uses sound waves to view the kidneys). The following tests may be appropriate: Doppler kidneys retroperitoneal ultrasound (which uses sound waves to view blood flow to and from the kidneys), abdominal and pelvic CT (<https://www.radiologyinfo.org/en/info/abdominect>) without contrast, MAG3 renal scan, abdomen MR angiography (<https://www.radiologyinfo.org/en/info/angiomr>) (MRI of the arteries) without contrast, abdominal and pelvic MRI (<https://www.radiologyinfo.org/en/info/mri-abdomen-pelvis>) without contrast, or abdominal MRI without contrast.

For chronic kidney disease, kidneys retroperitoneal ultrasound is usually appropriate. Abdominal and pelvic CT without contrast may be appropriate. Abdominal MR angiography without contrast, abdominal MRI without contrast, or Doppler kidneys retroperitoneal ultrasound may also be appropriate.

If doctors are unsure when kidney failure started, kidneys retroperitoneal ultrasound is usually appropriate. Abdominal and pelvic CT without contrast may be appropriate. Abdominal MR angiography without contrast, abdominal and pelvic MRI without contrast, or abdominal MRI without contrast may also be appropriate.

Kidneys retroperitoneal ultrasound is usually appropriate imaging for neurogenic bladder (the bladder may not function in the right way because of illness or injury). Tests that may be appropriate include abdominal and pelvic CT without contrast, DMSA renal scan (which gives detailed pictures of the kidneys), or abdominal and pelvic MRI without contrast.

— By Sandra V. Kotsis, MPH and Orit A. Glenn, MD. This information originally appeared in the *Journal of the American College of Radiology*.

Disclaimer

This information is copied from the RadiologyInfo Web site (<http://www.radiologyinfo.org>) which is dedicated to providing the highest quality information. To ensure that, each section is reviewed by a physician with expertise in the area presented. All information contained in the Web site is further reviewed by an ACR (American College of Radiology) - RSNA (Radiological Society of North America) committee, comprising physicians with expertise in several radiologic areas.

However, it is not possible to assure that this Web site contains complete, up-to-date information on any particular subject. Therefore, ACR and RSNA make no representations or warranties about the suitability of this information for use for any particular purpose. All information is provided "as is" without express or implied warranty.

Please visit the RadiologyInfo Web site at <http://www.radiologyinfo.org> to view or download the latest information.

Note: Images may be shown for illustrative purposes. Do not attempt to draw conclusions or make diagnoses by comparing these images to other medical images, particularly your own. Only qualified physicians should interpret images; the radiologist is the physician expert trained in medical imaging.

Copyright

This material is copyrighted by either the Radiological Society of North America (RSNA), 820 Jorie Boulevard, Oak Brook, IL 60523-2251 or the American College of Radiology (ACR), 1891 Preston White Drive, Reston, VA 20191-4397. Commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is prohibited.

